

King Fahd University of Petroleum and Minerals

Department of Mathematics and Statistics

Math 601

Comprehensive Exam– Term 232

Wednesday, January 24 , 2024

Allowed Time: 150 minutes

Name: _____

ID #: _____

Section #: _____

Serial Number: _____

Instructions:

1. Write clearly and legibly. You may lose points for messy work.
2. **Show all your work.** No points for answers without justifications !

Question #	Grade	Maximum Points
1		15
2		12
3		12
4		15
5		13
6		16
7		17
Total:		100

Exercise 1:(15)

Let X be a continuous random variable that takes only nonnegative values with mean μ and variance σ^2 .

A-(08) i)- Prove that for any value $a > 0$

$$P\{X \geq a\} \leq \frac{\mathbb{E}(X)}{a} \quad (\text{a})$$

ii)- Prove that for any $k > 0$

$$P(|X - \mu| \geq k) \leq \frac{\sigma^2}{k^2} \quad (\text{b})$$

Hint: You may use the result in (a)

B-(07) Suppose we know that the number of items produced in a factory during a week is a random variable with mean 500.

1- Find the probability that this week's production will be at least 1000 ?

2- If the variance of a week's production is equal 100, then find the probability that this week's production will be between 400 and 600 ?

Hint: You may use the results in A).

Exercise 2: (12)

Suppose the joint density of two random variables X and Y is given by:

$$f(x, y) = \begin{cases} 6xy(2 - x - y), & 0 < x < 1, 0 < y < 1 \\ 0, & \text{otherwise,} \end{cases} \quad (\text{c})$$

Compute the conditional expectation of X given that $Y = y$.

Exercise 3:(12)

The probability density function (pdf) of the duration of the (independent) interarrival times between successive cars on Dammam-Riyadh Highway is given by

$$f_T(t) = \begin{cases} \frac{1}{12} e^{-\frac{t}{12}}, & t \geq 0, \\ 0, & t < 0, \end{cases} \quad (\text{d})$$

where these durations are measured in seconds.

A-(04) An old fennec fox requires 12 seconds to cross the highway, and he starts out immediately after a car goes by. What is the probability that he will survive?

B-(04) Another old fennec, slower but tougher, requires 24 seconds to cross the road, but it takes two cars to kill him. If he starts out at an arbitrary time, determine the probability that he survives.

C-(04) If both these fennec foxes start out at the same time, immediately after a car goes by, what is the probability that exactly one of them survives?
(**Hint:** Consider a random variables N_1 = the number of cars in the first 12 seconds and N_2 = the number of cars in the second 12 seconds.)

Exercise 4: (15)

Let (Ω, \mathcal{F}, P) be a probability space equipped with filtration $(\mathcal{F}_t)_{t \geq 0}$ and $B_t, t \geq 0$ be the standard Brownian motion with respect to P and \mathcal{F}_t .

Show that the process Y defined by $Y_t := t^2 B_t^3, t \geq 0$, satisfies the stochastic differential equation:

$$dY_t = \left(\frac{2}{t} Y_t + 3(t^4 Y_t)^{\frac{1}{3}} \right) dt + 3(t Y_t)^{\frac{2}{3}} dB_t, \quad Y_0 = 0. \quad (\text{e})$$

Exercise 5:(13)

Let $B(t)$ be a 1-dimensional Brownian motion and $Y(t) = (Y_1(t), Y_2(t)) = (\cos(B(t)), \sin(B(t)))$. Show that the Itô process $Y(t)$ can be written in the following form:

$$dY_t = a Y_t dt + K Y_t dB_t, \quad (\text{f})$$

where a is a constant and K a suitable matrix.

Exercise 6:(16)

The charge $Q(t)$ at time t at a fixed point in an electric circuit satisfies the 2-dimensional stochastic differential equation:

$$\begin{cases} LQ_t'' + RQ_t' + \frac{1}{C} Q_t = G_t + \alpha W_t \\ Q(0) = Q_0, Q'(0) = I_0 \text{ given,} \end{cases} \quad (\text{g})$$

where W_t is a one-dimensional white noise, L, R, C, α, I_0 are constants and G_t a given function.

1-(08) Put

$$X_t(\cdot) = \begin{pmatrix} X_1 = Q_t \\ X_2 = Q_t' \end{pmatrix} \quad (\text{h})$$

and show that the SDE (g) can be written in the following form

$$dX_t = A X_t dt + H(t) dt + K dB_t, \quad (\text{i})$$

where A, H and K are suitable matrices and B_t a Brownian motion.

2-(08) Find the solution X_t of the stochastic differential equation (i).

Exercise 7:(17)

To model the spot freight rate in shipping, J.Tvedt(1995) used the geometric mean reversion process X_t which is defined as the solution of the stochastic differential equation

$$dX_t = \kappa (\alpha - \log X_t) X_t dt + \sigma X_t dB_t; \quad X_0 = x > 0, \quad (j)$$

where κ , α , σ and x are positive constants.

1-(08) Use the substitution $Y_t = \log X_t$ to transform the equation (j) into a linear stochastic differential equation for Y_t .

2-(06) Solve the Linear SDE obtained in 1). (**Hint:** Apply Itô formula to $e^{\kappa t} Y_t$).

3-(03) Find the solution X_t of the SDE (j).