

King Fahd University of Petroleum and Minerals
Department of Mathematics
Math 201
Final Exam
212
May 16, 2022
Net Time Allowed: 135 Minutes

MASTER VERSION

1. The absolute minimum of

$$f(x, y) = x^2 - y^2 + 4y$$

on the disc $x^2 + y^2 \leq 9$ is equal to

- (a) -21
- (b) 4
- (c) 11
- (d) 0
- (e) 3

(correct)

2. Let C be a curve given by the polar equation $r = \sec \theta(2 + \tan \theta)$.
The slope of the tangent line to the curve C at $\theta = \frac{\pi}{4}$ is

- (a) 4
- (b) 2π
- (c) π
- (d) 2
- (e) -2

(correct)

3. If M and m are the maximum and minimum values of

$$f(x, y) = xy$$

subject to $4x^2 + y^2 = 8$, then $M - m =$

- (a) 4
- (b) 0
- (c) 1
- (d) 2
- (e) -3

(correct)

4. The average value of

$$f(x, y) = x \sin y$$

over the region $R = \{(x, y) \mid 0 \leq x \leq 1, 0 \leq y \leq \pi\}$ is

- (a) $\frac{1}{\pi}$
- (b) 0
- (c) $\frac{-2}{\pi}$
- (d) $\frac{\pi}{2}$
- (e) $\frac{2}{\pi}$

(correct)

5. $\int_0^{\ln 2} \int_{e^y}^2 \frac{y^4 + 3}{x^2 + 1} dx dy =$

(a) $\int_1^2 \int_0^{\ln x} \frac{y^4 + 3}{x^2 + 1} dy dx$

(correct)

(b) $\int_0^2 \int_1^{\ln x} \frac{y^4 + 3}{x^2 + 1} dy dx$

(c) $\int_1^{\ln x} \int_1^2 \frac{y^4 + 3}{x^2 + 1} dy dx$

(d) $\int_1^2 \int_1^{\ln x} \frac{x^4 + 3}{y^2 + 1} dx dy$

(e) $\int_0^{\ln 2} \int_{e^x}^2 \frac{y^4 + 3}{x^2 + 1} dx dy$

6. The volume of the solid that lies under the paraboloid $z = x^2 + y^2$, above the xy -plane, and inside the cylinder $x^2 + y^2 = 2y$ is given by

(a) $\int_0^\pi \int_0^{2 \sin \theta} r^3 dr d\theta$

(correct)

(b) $\int_0^{2\pi} \int_0^{2 \sin \theta} r^3 dr d\theta$

(c) $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2 \cos \theta} r^3 dr d\theta$

(d) $\int_0^\pi \int_0^{\sin \theta} r^2 dr d\theta$

(e) $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2 \cos \theta} r^2 dr d\theta$

7. The volume of the solid bounded by $z = x^2 + y^2$, and $z = 2 - x^2 - y^2$ is equal to

(a) π

(correct)

(b) $\frac{\pi}{2}$

(c) 3π

(d) 4π

(e) $\frac{\pi}{4}$

8. If $E = \{(x, y, z) \mid 0 \leq x \leq 1, 0 \leq y \leq 1, 0 \leq z \leq 1\}$, then $\int \int \int_E xyz^2 dV =$

(a) $\frac{1}{12}$

(correct)

(b) $\frac{1}{26}$

(c) $\frac{1}{3}$

(d) $\frac{1}{2}$

(e) $-\frac{1}{6}$

9. The volume of the tetrahedron T bounded by the planes $x + 2y + z = 2$, $x = 2y$, $x = 0$ and the xy -plane is equal to

(a) $\frac{1}{3}$

(correct)

(b) $\frac{2}{3}$

(c) $\frac{4}{3}$

(d) $\frac{1}{2}$

(e) $\frac{1}{6}$

10. The volume of the solid bounded by $x^2 + (y - 1)^2 = 1$, and the planes $z = 0$ and $x + z = 2$ is given by

(a) $\int_0^\pi \int_0^{2 \sin \theta} \int_0^{2-r \cos \theta} r \, dzdrd\theta$

(correct)

(b) $\int_0^{2\pi} \int_0^{2 \cos \theta} \int_0^{1-r \cos \theta} dzdrd\theta$

(c) $\int_0^{\frac{\pi}{2}} \int_0^{2 \sin \theta} \int_0^{1-r \cos \theta} dzdrd\theta$

(d) $\int_0^\pi \int_0^{2 \sin \theta} \int_0^{1-r \cos \theta} dzdrd\theta$

(e) $\int_0^{2\pi} \int_0^{2 \cos \theta} \int_0^2 r \, dzdrd\theta$

11. The volume of the solid E that lies above the cone $\phi = \frac{\pi}{6}$ and below the sphere $\rho = 2 \cos \phi$ is

(a) $\frac{7\pi}{12}$

(correct)

(b) $\frac{\pi}{3}$

(c) $\frac{\pi}{4}$

(d) $\frac{5\pi}{11}$

(e) $\frac{3\pi}{4}$

12. $\int_0^{2\pi} \int_0^\pi \int_0^{\frac{1-\cos\phi}{2}} \rho^2 \sin\phi \, d\rho d\phi d\theta =$

(a) $\frac{\pi}{3}$

(correct)

(b) $\frac{\pi}{2}$

(c) $\frac{\pi}{7}$

(d) $\frac{5\pi}{4}$

(e) π

13. The area of the region enclosed by one loop of the curve $r = \sin 3\theta$ is

(a) $\frac{\pi}{12}$

(correct)

(b) $\frac{\pi}{2}$

(c) $\frac{\pi}{7}$

(d) $\frac{5\pi}{3}$

(e) 2π

14. $\lim_{(x,y) \rightarrow (1,1)} \frac{xy^2 - 1}{y - 1} =$

(a) does not exist

(correct)

(b) 0

(c) -1

(d) 1

(e) ∞

15. A point on the cone $x^2 + y^2 - z^2 = 0$, where the tangent plane is parallel to the plane $3x + 4y + 5z = 0$ is

(a) $(3, 4, -5)$

(correct)

(b) $(3, 1, 1)$

(c) $(2, 4, 5)$

(d) $(2, 3, 3)$

(e) $(1, 1, -5)$

16. The volume of the solid bounded by the cylinder $x^2 + y^2 - 2y = 0$ on the lateral sides and bounded on top and bottom by the sphere $x^2 + y^2 + z^2 = 4$ is given by

(a) $\int_0^\pi \int_0^{2\sin\theta} \int_{-\sqrt{4-r^2}}^{\sqrt{4-r^2}} r \, dzdrd\theta$

(correct)

(b) $\int_0^{\frac{\pi}{2}} \int_0^{2\sin\theta} \int_{-\sqrt{2-r^2}}^{\sqrt{2-r^2}} r \, dzdrd\theta$

(c) $\int_0^{\frac{\pi}{2}} \int_0^{\sin\theta} \int_{-\sqrt{4-r^2}}^{\sqrt{4-r^2}} dzdrd\theta$

(d) $\int_0^\pi \int_0^{2\cos\theta} \int_0^{\sqrt{4-r^2}} r \, dzdrd\theta$

(e) $\int_0^\pi \int_0^{2\cos\theta} \int_{-\sqrt{4-r^2}}^{\sqrt{4-r^2}} dzdrd\theta$

17. The distance from the point $P(5, 6, -1)$ to the line

$$L : x = 2 + 8t, \quad y = 4 + 5t, \quad z = -3 + 6t$$

is equal to

(a) $\frac{3}{5\sqrt{5}}$

(correct)

(b) $\frac{1}{\sqrt{5}}$

(c) $\frac{3}{\sqrt{5}}$

(d) $\frac{2}{\sqrt{5}}$

(e) $\frac{4}{3\sqrt{5}}$

18. The normal line of the surface $4x^2 + y^2 + 2z = 6$ at the point $P(-1, 2, -1)$ passes through the point

(a) $(-9, 6, 1)$

(correct)

(b) $(7, 2, 1)$

(c) $(5, -6, 1)$

(d) $(3, 3, 2)$

(e) $(9, 9, 6)$

19. Let C be a curve given by the parametric equations

$$x = \frac{1}{t}, \quad y = t - \sin(\pi t), \quad t > 0.$$

The slope of the tangent line to the curve C at $t = 1$ is

(a) $-1 - \pi$

(correct)

(b) $2 - \pi$

(c) $2 + \pi$

(d) π

(e) $-2 + 2\pi$

20. The function

$$f(x, y) = x^3 + y^3 - 6xy$$

has

(a) one local minimum

(correct)

(b) one local maximum

(c) two saddle points

(d) two local maxima

(e) no saddle points

21. If $z = \tan^{-1}\left(\frac{u^2}{\sqrt{v}}\right)$, where $u = 2y - x$ and $v = 3x - y$.

Then $\frac{\partial z}{\partial y}$ at $(x, y) = (2, 2)$ is

(a) $\frac{17}{20}$

(b) $\frac{7}{2}$

(c) $\frac{14}{5}$

(d) 2

(e) $\frac{2}{3}$

(correct)